

HOW OZEMPIC® WORKS



Keeps your blood sugar in check^{1,2}



Helps you manage your weight^{3,4}



Protects your heart & kidneys^{5,6}



OZEMPIC®
semaglutide injection

When to inject Ozempic®

The starting dose of Ozempic® is 0.25 mg.⁶

Ozempic® should be dosed **once a week on the same day each week.**




The injection can be dosed at **any time** of the day.




How to use Ozempic® FlexTouch®⁶

Ozempic® FlexTouch® is simple to use!




Subcutaneous Injection under the skin



FlexTouch® demo video

Scan the QR code to watch the FlexTouch® video



100 x 65 mm

100 x 65 mm

100 x 65 mm

100 x 65 mm

100 x 65 mm

How to store Ozempic® FlexTouch®⁶

Before First Use



2°C to 8°C

After First Use

For 6 weeks



Below 30°C

Note: Do not freeze and do not use if it has been frozen. Keep away from sunlight.

100 x 65 mm

How to travel with Ozempic® FlexTouch®^{6,7}

Ozempic® FlexTouch® is convenient to carry!

Carry in **cabin baggage or a handbag** and avoid freezing



Avoid storing in a **car or glove box**, as heat may lower its potency



100 x 65 mm

Missed Dose⁶

Missed a dose?
Here's what to do next!




Diagram illustrating the missed dose protocol:

- Day 1: Missed Dose (X)
- Day 2: Take Missed Dose (✓)
- Day 3: Take Missed Dose (✓)
- Day 4: Take Missed Dose (✓)
- Day 5: Take Missed Dose (✓)
- Day 6: Do Not Take Missed Dose (X)
- Day 7: Do Not Take Missed Dose (X)
- Day 8: Take Next Dose (✓)

100 x 65 mm

Well-being during treatment^{6,8}

If you notice any side effects, don't worry - they're usually temporary. Here are a few simple tips to help prevent or ease them.

Eat smaller meals



Stop eating when full or when not hungry



Avoid high-fat, fried, and spicy foods



Avoid carbonated and alcoholic beverages



Drink plenty of water and eat high-fibre foods



100 x 65 mm

ONCE-WEEKLY

OZEMPIC®

semaglutide injection

Reference: 1. Aroda VR et al. Diabetes & metabolism. 2019 Oct 1;45(5):409-18. | 2. Pratley R E et al., Presented at the 78th Scientific Sessions of the American Diabetes Association, 22–26 June 2018, Orlando, FL, USA. | 3. Åhrén B et al. Diabetes, Obesity and Metabolism. 2018 Sep;20(9):2210-9. | 4. Marso SP et al. New England Journal of Medicine. 2016 Nov 10;375(19):1834-44. | 5. Perkovic V et al. New England Journal of Medicine. 2024 Jul 11;391(2):109-21. | 6. Ozempic® CDSCO approved package insert version dated 23 Oct 2025. | 7. Adopted from <https://www.medscape.com/viewarticle/what-your-patients-should-know-about-traveling-glp-1s-2025a1000mmi?form=fpf> last accessed on 17/11/25. | 8. Gorgojo-Martínez JJ, et al. Journal of clinical medicine.2022 Dec 24;12(1):145.


This does not replace medical advice. Talk to your doctor/ health care professional for more details.
The Apis bull logo is a registered trademarks owned by Novo Nordisk A/S and registered in Denmark. Novo Nordisk India Private Limited, Nxt Tower-2, Floor 1 & 2, Embassy Manyata Business Park, Nagavara Village, Kasaba Hobli, Bangalore-560 045, India



IN25OZM00040 | Review Date: 19 Nov 2025



100 x 65 mm



Closed size 105 x 70 mm